

Percentage of susceptible Organisms Isolated From Stool and Rectal Swab, Clinical Microbiology Laboratory, Vachiraphuket Hospital, January-December-2023

Organism	TOTAL ISOLATES	BETA - LACTAMS									CARBAPENEMS		QUINOLONES		AMINOGLYCOSIDES		MISCELLANEOUS		
		AMPICILLIN	AMOXICILLIN/CLAVULANIC ACID	AMPICILLIN /SULBACTAM	PIPERACILLIN /TAZOBACTAM	CEFOTAXIME	CEFTAZIDIME	CEFTRIAXONE	CEFEPIME	CEFOXITIN	IMIPENEM	MEROPENEM	CIPROFLOXACIN	LEVOFLOXACIN	AMIKACIN	GENTAMICIN	CHLORAMPHENICOL	CO-TRIMOXAZOLE	TETRACYCLINE
<i>Salmonella</i> , Non-typhoidal	79	77 (79)						100 (79)					90 (79)					95 (79)	
<i>Vibrio parahaemolyticus</i>	5 ^a	0 (5)						100 (5)					100 (5)					100 (5)	

a : Calculated from fewer than the standard recommendation of 30 isolates

^b : Blood, Pleural Fluid, CSF

^c : Sputum, Ear, Sinus

^d : BY MIC (E-test)

^e : MIC Interpretive Criteria

^f : Interpret according to oxacillin MICs susceptibility test

^g : COLISTIN จำนวนจาก Intermediate COLISTIN BY MIC (Microbroth dilution method)

^h : High-Level Aminoglycoside ;Susceptible: is synergistic with cell wall-active agent (eg. ampicilin, penicillin and

^u : Urine, Urine Catheter, Urine Clean- Voided

R : Intrinsic resistance

