

Percentage of susceptible Organisms Isolated From Stool and Rectal Swab, Clinical Microbiology Laboratory, Vachiraphuket Hospital, January-December-2021

Organism	TOTAL ISOLATES	BETA - LACTAMS											CARBAPENEMS		QUINOLONES		AMINOGLYCOSIDES		MISCELLANEOUS		
		AMPICILLIN	AMOXICILLIN/ CLAVULANIC ACID	AMPICILLIN / SULBACTAM	PIPERACILLIN / TAZOBACTAM	CEFAZOLIN (A)	CEFUROXIME SODIUM (parenteral)	CEFOTAXIME	CEFTAZIDIME	CEFEPIME	CEFOXITIN	IMIPENEM	MEROPENEM	CIPROFLOXACIN	LEVOFLOXACIN	AMIKACIN	GENTAMICIN	CHLORAMPHENICOL	CO-TRIMOXAZOLE	TETRACYCLINE	
<i>Salmonella</i> , Non-typhoidal	79	68 (79)											86 (79)					92 (79)			
<i>Vibrio parahaemolyticus</i>	6 ^a	20 (6)						100 (6)					100 (6)					100 (6)			

^a : Calculated from fewer than the standard recommendation of 30 isolates

^b : Blood, Pleural Fluid, CSF

^c : Sputum, Ear, Sinus

^d : Interpret according to oxacillin susceptibility test

^e : MIC Interpretive Criteria

^f : Interpret according to cefoxitin susceptibility test

^u : Urine, Urine Catheter, Urine Clean- Voided

R : Intrinsic resistance

^h : High-Level Aminoglycoside ;Susceptible: is synergistic with cell wall-active agent (eg. ampicilin, penicillin and vancomycin).

